

CARE Clinic

Cranial Asymmetry Repositioning Evaluation (CARE)

For infants with plagiocephaly and torticollis, we offer an enhancement to our helmet assessments, with the addition of a physical therapist to address further needs, all in one appointment.

Infant Plagiocephaly and Torticollis Diagnoses

These diagnoses are on the rise and often occur together. LifeScape has responded to this with the CARE (Cranial Asymmetry Repositioning Evaluation) Clinic, to screen for and treat both conditions. The clinic's team of physical therapists and orthotists aim to provide comprehensive CARE for this complex diagnosis.

Early Detection

For both torticollis and plagiocephaly, early detection is critical for effective treatment and parent education. This is because both can have an affect on a child's development and facial alignment. Evaluations for both conditions are available at LifeScape in one convenient appointment.



For more information:

- **Call** 712.226.2253 for details on the CARE Clinic at LifeScape Pediatric Therapy in Sioux City
- **Location** 2524 Glenn Avenue
- **Web** RiseCustomSolutions.com




The Most Advanced Technology – to Deliver the Best Results

LifeScape Pediatric Therapy in Sioux City uses The SmartSoc® 3D Capturing System, which is the latest addition to Orthomerica's US FDA 510(k) cleared line of image capture devices for manufacturing of STARband® and STARlight® Cranial Orthoses. The system is portable and provides quick and accurate images – keeping the baby happy.


The CARE Clinic provides a team approach for the treatment of plagiocephaly and torticollis.

CARE Clinic

Cranial Remolding Evaluation & Treatment



Before Treatment Scan



After Treatment Scan



Before STARband® Treatment



After STARband® Treatment



For more information:

- **Call** 712.226.2253 for details on the CARE Clinic at LifeScape Pediatric Therapy in Sioux City
- **Location** 2524 Glenn Avenue
- **Web** RiseCustomSolutions.com

Torticollis Evaluation & Treatment



Infant Before Treatment



Infant After 4 Months of Treatment